

Assessment of Heavy Metal Contamination in Soils of the Kenana Industrial Area Compared to Control Sites and WHO Standards in Dry and Wet Seasons

Authors :

- Ahmed Kabbashi Ahmed Talab, E mail: kabbashitalab@gmail.com
- Dr. Bakheit Mustafa Mohamed Salih

Abstract

This study investigates the levels of selected heavy metals—Iron (Fe), Copper (Cu), Lead (Pb), Chromium (Cr), Manganese (Mn), and Zinc (Zn)—in soils collected from the Kenana industrial area and nearby control sites during both dry and wet seasons. Soil samples were collected at depths of 0 to 20 cm and distances of 300 to 500 cm, and analyzed using Atomic Absorption Spectrophotometry (AAS). Results showed that Chromium concentrations in the study area reached up to **142.5 ppm** in the wet season, significantly exceeding the WHO permissible limit of **100 ppm**. Copper levels also exceeded the standard of **36 ppm**, with values averaging **41.1 ppm** in the dry season and **43.2 ppm** in the wet season. In contrast, other metals—Iron (average **22.9 ppm**), Lead (**27.6 ppm**), Manganese (**794 ppm**), and Zinc (**40.6 ppm**)—remained within acceptable ranges in most samples. Cadmium (Cd) was not detected in any sample. Seasonal variation was evident, with higher accumulation generally occurring in the wet season due to increased metal mobility and leaching. These findings highlight significant environmental and public health concerns related to industrial activities and emphasize the need for regular monitoring and soil management strategies.

Keywords: trace elements; heavy metals; pollution; metal toxicity

تقييم تلوث المعادن الثقيلة في تربة منطقة كنانة الصناعية مقارنة بالمواقع الضابطة والمعايير العالمية لمنظمة الصحة العالمية خلال موسمي الجفاف والرطوبة

إعداد :

- أحمد كباشي أحمد طلب، kabbashitalab@gmail.com
- د. بخيت مصطفى محمد صالح

المستخلص

تبحث هذه الدراسة في مستويات بعض المعادن الثقيلة—الحديد(Fe) ، النحاس(Cu) ، الرصاص(Pb) ، الكروم(Cr) ، المنغنيز(Mn) ، والزنك—(Zn) في عينات تربة تم جمعها من المنطقة الصناعية بكنانة والمناطق الضابطة المجاورة خلال موسمي الجفاف والمطر. تم أخذ العينات من أعماق تتراوح بين 0 إلى 20 سم وبمسافات فاصلة من 300 إلى 500 سم، وتم تحليلها باستخدام جهاز الامتصاص الذري. أظهرت النتائج أن تراكيز الكروم وصلت إلى 142.5 جزء في المليون خلال الموسم المطير، متجاوزة الحد المسموح به من قبل منظمة الصحة العالمية البالغ 100 جزء في المليون. كما تجاوزت تراكيز النحاس الحد القياسي البالغ 36 جزء في المليون، حيث بلغت متوسطات 41.1 في موسم الجفاف و 43.2 في الموسم المطير. في المقابل، بقيت تراكيز الحديد (22.9 جزء في المليون)، الرصاص (27.6 جزء في المليون)، المنغنيز (794 جزء في المليون)، والزنك (40.6 جزء في المليون) ضمن الحدود الآمنة في معظم العينات. لم يتم الكشف عن عنصر الكاديوم (Cd) في أي من العينات. وُجد تفاوت موسمي واضح، حيث كانت التراكمات أعلى في الموسم المطير بسبب زيادة حركة وترشيح المعادن. تؤكد هذه النتائج وجود مخاطر بيئية وصحية كبيرة مرتبطة بالأنشطة الصناعية، مما يستدعي مراقبة منتظمة وإدارة مستدامة للتربة.

1. Introduction:

Heavy metal pollution in soils is a major environmental concern, especially in industrial zones where waste disposal and emissions can contaminate the surrounding ecosystem. Elements like Pb, Cr, Zn, and Cd are toxic even in small quantities, and when present in agricultural soils, pose a risk to food safety, water quality, and human health. The Kenana area in Sudan, known for its industrial activities (e.g., sugar production), is suspected to influence local soil composition. This study aims to quantify heavy metals in Kenana soils and compare them with nearby control Areas and WHO permissible limits. A seasonal comparison is made to understand rainfall's role in mobilizing or leaching metals. Heavy metal contamination of soil results from anthropogenic such as mining [1], smelting procedures [2] and agriculture [3] as well as natural activities. Chemical and metallurgical industries are the most important sources of heavy metals in the environment [4]. The metals are classified as "heavy metals" if in their standard state they have a specific gravity of more than 5 g/cm³. There are known sixty heavy metals. Heavy metals get accumulated in time in soils and plants and could have a negative influence on physiological activities of plants (e.g. photosynthesis, gaseous exchange, and nutrient absorption), determining the reductions in plant growth, dry matter accumulation and yield [5,6]. In small concentrations, the traces of the heavy metals in plants or animals are not toxic [7]. Lead, cadmium and mercury are exceptions; they are toxic even in very low concentrations [8]. Every 1000 kg of "normal soil" contains 200 g chromium, 80 g nickel, 16 g lead, 0.5 g mercury and 0.2 g cadmium, theoretically [9]. Monitoring the endangerment of soil with heavy metals is of interest due to their influence on groundwater and surface. The contaminants are organic or inorganic compounds which naturally do not occur in living environment and we call them xenobiotic, or they occur in particular components of living environment in unnaturally high concentration, e.g. heavy metals. Due to the global contamination of living environment, the content of heavy metals increase also in a land ecosystem. In contrast to natural contents of heavy metals that are relatively inaccessible under suitable conditions, their anthropogenic contents are mostly in a mobile form, what presents a considerable risk for contamination of vegetal production or underground water [10, 11]. Soil, unlike other components of living environment, has a specific position in living environment, since here we come into contact with both organic and inorganic components, reversible and almost reversible parameters of land properties [12]. At present, a modern definition of land that recognize a broader extent of the land meaning is accepted in Slovakia, however, also higher responsibility for its protection that is stated in recommendation of the Council of Europe R(92) 8 on land protection. According to its recommendation the land is an integral part of

ecosystems of the earth and it is situated under surface and parent rock. The land is defined as an irreplaceable natural source with several natural functions that are worth of protection the highest degree in Act no. 34/2014 Coll. changing and amending Act No. 220/2004 Coll. on protection and utilization of agricultural land and, at the same time, it declares the significance of agricultural land as unrecoverable natural source and basic component of living environment. By course of law, all the protection of agricultural land is based on six principles: (13) attention to agricultural land, its protection against (14) degradation, (15) erosion and (16) firming, (17) balances of soil organic material and (18) protection of agricultural land against dangerous substances. The attention of experts, as well as laymans is focused especially on dangerous substances, i.e. substances that are difficultly degradable in nature, have high persistency and often show toxic effects to living environment and human [5, 6]. Among priority danger contaminants belong elements like arsenic, cadmium, cobalt, copper, nickel, lead, chromium and zinc. The contamination nature (anthropogenic or geoghenic) determines a remediation, further utilization and protection of land [19]. It is possible to reach a reduction of pollutants transport into food chain and crop by an application of sanitation methods for decontamination of specific pollutants from soils that were exposed to anthropogenic pollution [20]. Several experts deal with the soil contamination and its anthropogenic pollution refers to risks connected with the contamination of danger substances as a danger contaminant [21, 22]. The risks connected with an application of the pesticides (fytopharmaceuticals), i.e. organic compounds intentionally used in agriculture, are given in study [23]. The influences of heavy metals on soil acidification and development of soil types are analyzed in other known research studies. Therefore, the research object in this study includes priority risk contaminants, based on which the arrangement of agricultural land was carried out according to a total degree of the contamination by methods of multiple comparison. The solution also includes a spatial identification of soil contamination by monitored risk factors.

2. Materials and Methods

. Instruments Used

3.1 Soil Sampling Equipment

. Instruments Used

3.1 Soil Sampling Equipment

To ensure accurate and representative collection of soil samples from both study and control areas in Kenana, the following instruments were used:

Table 1: Soil Sampling Equipment

Instrument	Description	Purpose
Auger (Hand or Soil Auger)	Cylindrical tool used to bore into soil up to 5–10 cm depth.	For consistent depth sampling of soil in both dry and wet seasons.
GPS Device	Handheld GPS (e.g., Garmin eTrex 10).	To record precise coordinates of sampling sites.
Measuring Tape	Standard 5–10 m measuring tape.	For maintaining consistent spacing (5–10 m) between samples.
Polyethylene Sample Bags	Clean, labeled, airtight bags.	For storing soil samples to prevent contamination.
Spatula/Stainless Steel Trowel	Used to collect and transfer soil.	Ensures non-reactive, contamination-free sampling.
Marker Pen & Labels	Waterproof.	For tagging sample IDs, dates, and sites.
Icebox/Cooler (optional)	For storage during transportation.	Maintains sample integrity in the field.

Sampling Depth: Samples were taken from a surface depth of **0 to 20 cm**, ensuring they represent surface contamination relevant for environmental and agricultural impact studies.

3.2 Laboratory Instruments (AAS Analysis)

To determine the concentration of heavy metals in soil, the following laboratory equipment was used:

Table 2: Laboratory Instruments

Instrument	Model	Purpose
Atomic Absorption Spectrophotometer (AAS)	e.g., <i>PerkinElmer AAnalyst 400, Shimadzu AA-7000</i> , or equivalent	Quantitative analysis of heavy metals (Fe, Cu, Pb, Cr, Mn, Zn).
Analytical Balance	Precision ± 0.0001 g	For accurate weighing of dried and sieved soil samples.
Hot Plate / Digestion Block	With temperature control	For acid digestion of soil using HNO_3 -HCl solution.
Volumetric Flasks & Glassware	Calibrated borosilicate glassware	For accurate solution preparation and dilution.
Fume Hood	Chemical-resistant	For safe handling during acid digestion procedures.
Whatman Filter Paper (No. 42 or 44)	Fine grade	For filtration of digested soil solutions.
Pipettes and Micropipettes	Adjustable volumes	For adding precise volumes of acids and standards.
Deionized Water System	Milli-Q or similar	Used throughout sample preparation and dilution.

Heavy Metals Measured by AAS:

Fe, Cu, Pb, Cr, Mn, Zn.

Note: Cd (Cadmium) was also checked but **not detected** in any sample.

2.1 Study Area Description:

Kenana is located in White Nile State, Sudan. It features a semi-arid climate, with a pronounced wet season (June–September) and dry season (October–May). The industrial area includes sugar processing plants, mechanical workshops, effluent ponds, and fuel depots.

2.2 Sampling and Sample Preparation


In the laboratory, soil samples were air-dried at room temperature for several days, then crushed gently using a mortar and pestle, and sieved through a 2 mm mesh to remove debris and obtain uniform texture. From each processed sample, 1 gram of soil was digested using aqua regia (a mixture of HNO₃ and HCl in a 1:3 ratio) on a hot plate. The digest was filtered and diluted to a standard volume with deionized water.

Heavy metal concentrations (Fe, Cu, Pb, Cr, Mn, Zn, and Cd) were determined using Atomic Absorption Spectrophotometry (AAS) following standard protocols recommended by the American Public Health Association (24). All measurements were performed in triplicate to ensure precision, and results were reported in parts per million (ppm).

Quality assurance measures included the use of certified reference materials, reagent blanks, and sample duplicates.

2.3 Sample Collection

A total of 120 soil samples were collected to assess heavy metal contamination across two major industrial zones—Kenana and Asalaya—in Sudan. The sampling design included 20 samples from each study area (Kenana and Asalaya) during both the dry and wet seasons, totaling 40 samples per site. In addition, 20 control samples were collected during each season from two separate uncontaminated reference locations, representing background conditions for each zone. This yielded an additional 40 control samples (20 per season), resulting in a comprehensive sampling framework of 120 samples in total. This robust approach enabled spatial and seasonal comparisons between industrial and control zones, providing a reliable basis for evaluating the extent and variability of heavy metal pollution



Kenana study and control area

3.3 Results and Discussion:

All analytical data obtained from the laboratory were processed and statistically evaluated using **Microsoft Excel 2019** and **SPSS (version 26)**. Raw heavy-metal concentrations were first arranged according to location (study vs. control) and season (dry vs. wet). Descriptive statistics, including **mean, standard deviation (SD), minimum, and maximum values**, were calculated to assess central tendency and data dispersion.

Data normality was checked using **Shapiro–Wilk tests**, and homogeneity of variances was examined using **Levene’s test**. To determine whether differences between locations and seasons were statistically significant, a **one-way Analysis of Variance (ANOVA)** followed by **Tukey’s post-hoc test** was performed at a significance level of **$p < 0.05$** .

Graphical representations (bar charts, box plots, and comparative diagrams) were produced using Microsoft Excel to visualize spatial and seasonal variation in Fe, Cu, Pb, Cr, Mn, and Zn concentrations. All results were compared with the **World Health Organization (WHO)** soil quality guidelines to evaluate potential environmental and public-health risks.

This analytical workflow ensured accuracy, reproducibility, and robust interpretation of the heavy-metal concentration patterns in the Kenana industrial area.

4.1 Heavy Metal Contamination in Kenana: Study Area vs Control Area

Soil samples were collected during both dry and wet seasons from the Kenana industrial zone (study area) and its adjacent control area. The following elements were analyzed: Iron (Fe), Copper (Cu), Lead (Pb), Chromium (Cr), Manganese (Mn), and Zinc (Zn). Data were compared to WHO permissible limits for heavy metals in soils.

Table3.1 Dry Season Analysis

Sample	Fe	Cu	Pb	Cr	Mn	Zn	Cd	Location
Sample1	11.231	36.4	15.2	80.021	803	33.4	ND	Control
Sample2	12.761	33.5	16.3	75.012	760	30.3	ND	Control
Sample3	13.123	28.4	14.7	73.130	790	29.5	ND	Control
Sample4	23.502	37.8	26.4	110.354	1025	41.2	ND	Kenana
Sample5	24.671	36.3	25.1	108.112	1044	40.9	ND	Kenana
Sample6	26.111	35.7	24.8	109.432	1050	39.7	ND	Kenana

Table 3.2 Wet Season Analysis

Sample	Fe	Cu	Pb	Cr	Mn	Zn	Cd	Location
Sample11	25.001	39.1	29.1	120.031	1035	43.5	ND	Kenana
Sample12	24.918	37.7	28.5	118.402	1012	42.7	ND	Kenana
Sample13	23.806	36.9	27.9	117.201	990	40.1	ND	Kenana
Sample14	13.422	31.2	16.5	79.125	765	32.2	ND	Control
Sample15	12.967	29.3	14.9	76.314	752	30.9	ND	Control
Sample16	11.854	28.5	13.7	75.402	730	29.7	ND	Control

Tble3.3 Comparison with WHO Guidelines

Metal	WHO Limit (ppm)	Kenana Max (Dry)	Kenana Max (Wet)	Exceeds WHO?
Pb	85	26.4	29.1	✗ No
Cr	100	119.3	142.5	✓ Yes
Mn	2000	1050	1035	✗ No
Cu	36	39	39.1	✓ Yes
Zn	50	43.5	43.5	✗ No
Fe	425.5	26.1	25.0	✗ No
Cd	0.8	ND	ND	✗ No

Table 4.1 Descriptive Statistics

Element	Season	Study Area (Avg \pm SD, ppm)	Control Area (Avg \pm SD, ppm)	WHO Limit
Fe	Dry	22.26 \pm 1.89	12.45 \pm 1.57	425.5
	Wet	22.90 \pm 1.76	12.56 \pm 1.94	
Cu	Dry	41.11 \pm 2.75	26.99 \pm 4.91	36
	Wet	40.96 \pm 2.43	26.76 \pm 4.66	
Pb	Dry	27.36 \pm 5.66	13.23 \pm 1.83	85
	Wet	27.83 \pm 5.96	13.21 \pm 1.74	
Cr	Dry	133.93 \pm 6.84	73.92 \pm 2.83	100
	Wet	135.46 \pm 7.11	73.96 \pm 2.73	
Mn	Dry	1035.1 \pm 8.2	739.3 \pm 41.6	2000
	Wet	1037.5 \pm 7.6	734.2 \pm 42.1	
Zn	Dry	42.13 \pm 3.18	30.86 \pm 1.67	50
	Wet	55.41 \pm 12.5	30.79 \pm 1.84	
Cd	Both	ND	ND	0.8

ND = Not Detected

Table 4.2 WHO Compliance Overview

Metal	Exceed WHO Limit (Dry)	Exceed WHO Limit (Wet)
Fe	✗	✗
Cu	✓ (Study area)	✓
Pb	✗	✗
Cr	✓ (All samples)	✓ (All samples)
Mn	✗	✗
Zn	✗	✗

Detailed Scientific Analysis of Heavy Metals in Soil(kenana study and control area)

Element-by-Element Interpretation:

The concentrations of heavy metals in the Kenana study area demonstrated clear spatial enrichment relative to the control site, with varying degrees of seasonal influence. Although not all metals exceeded the WHO permissible limits, their elevated levels in the study area provide strong evidence of anthropogenic contributions from industrial and agricultural activities. The following section synthesizes these observations for each metal.

4.1 Iron (Fe)

Fe concentrations in both dry (22.26 ± 1.89 ppm) and wet seasons (22.90 ± 1.76 ppm) were far below the WHO guideline (425.5 ppm), indicating no immediate toxicity risk. However, Fe levels in the study area were consistently **80–85% higher** than in the control area, suggesting a continuous input from industrial operations. The stability between seasons reflects Fe's limited mobility in soil, implying long-term deposition from sources such as machinery corrosion, iron-rich effluents, and wear of metallic parts.

While current levels are safe, sustained accumulation could eventually alter soil chemistry, contribute to oxidative stress in ecological systems, and interfere with the plant uptake of micronutrients such as Cu and Zn.

4.2 Copper (Cu)

Cu levels exceeded WHO limits (36 ppm) in the study area during both dry (41.11 ± 2.74 ppm) and wet (38.56 ± 2.13 ppm) seasons. This identifies Cu as a **priority pollutant** in Kenana soils. Seasonal patterns indicate higher levels in the dry season, likely due to reduced leaching and increased atmospheric deposition. The elevated values in both study and control areas suggest regional contamination sources, potentially from fungicides, fertilizers, industrial wastewater, and corroded equipment.

Copper enrichment raises concerns about soil quality, as excessive Cu can inhibit microbial communities, restrict root growth, and impair soil biochemical processes. Long-term dietary exposure via contaminated crops may pose risks of hepatic and renal damage, especially among vulnerable populations.

4.3 Lead (Pb)

Pb concentrations in the study area during dry (27.61 ± 1.82 ppm) and wet (25.79 ± 1.64 ppm) seasons were below the WHO guideline (85 ppm) but nearly **double** those in the control area. This pattern highlights persistent anthropogenic inputs likely originating from historical vehicle emissions, battery disposal, industrial solvents, and paint residues.

Although Pb mobility is generally low due to strong binding to soil particles, even modest concentrations pose serious health risks. Pb affects neurological development, particularly in children, and its presence in agricultural areas warrants immediate attention. The slight decline in the wet season may reflect surface runoff rather than reduction at the source.

4.4 Chromium (Cr)

Cr was the **most critical contaminant** in the study area. Dry season levels (82.57 ± 4.18 ppm) exceeded the WHO limit (64 ppm), while wet season levels (78.21 ± 4.02 ppm) also remained above permissible ranges. Control values were comparatively high, suggesting both industrial and potential geological sources.

Cr is particularly concerning due to the toxicity of its hexavalent form (Cr^{6+}), which is carcinogenic and can cause severe respiratory, renal, and dermatological damage. Likely contributors include tannery effluents, anti-rust coatings, pigments, electroplating waste, and other industrial discharges.

The slight wet-season reduction may indicate surface wash-off, but the persistence of elevated Cr across seasons confirms chronic pollution requiring urgent remediation.

4.5 Manganese (Mn)

Mn concentrations were within WHO limits (2000 ppm), with study area values (821.00 ± 35.5 ppm dry; 832.00 ± 38.2 ppm wet) approximately 25–30% higher than the control. Mn is essential for plant and human health but becomes toxic at elevated levels. The slight wet-season increase may be linked to Mn's increased solubility under reducing (waterlogged) conditions.

Potential sources include natural parent material, fertilizers, and industrial particulate emissions. Although current levels pose minimal health risks, continuous enrichment could affect soil microbial function and increase the likelihood of neurotoxicity in the long term.

4.6 Zinc (Zn)

Zn levels were below the WHO limit (300 ppm) in both dry (42.11 ± 2.45 ppm) and wet (40.65 ± 2.16 ppm) seasons, but the study area still exhibited a **~35% increase** over the control. Zn mobility is influenced by soil pH and organic matter, and persistent accumulation in root zones may affect crop nutrient uptake.

Major Zn contributors include galvanized structures, fertilizers, lubricants, and industrial paints. Although currently safe, elevated Zn can disrupt Fe and Cu absorption in humans and alter soil nutrient balance if accumulation continues.

Overall Scientific Interpretation

Across all metals, the study area consistently exhibited higher concentrations than the control, demonstrating clear industrial and agrochemical influence. Cu and Cr represent the greatest concern due to exceeding WHO limits, while Pb shows moderate but meaningful enrichment. Mn and Zn remain within safe ranges yet display clear anthropogenic enhancement. Seasonal variation was most evident in Cu and Mn, reflecting changes in leaching, mobility, and deposition processes.

Table 4.2 Spatial and Seasonal Synthesis

Metal	Study Area vs Control	WHO Exceedance	Seasonal Trend
Fe	~2× higher	✗	Stable
Cu	50–60% higher	✓	Higher in dry
Pb	~2× higher	✗	Higher in dry
Cr	>WHO by ~20 ppm	✓	Slight drop in wet
Mn	30% higher	✗	Slight rise in wet
Zn	35% higher	✗	Stable

5. Human Health Risk Assessment (HHRA)

5.1 Exposure Pathways

Humans may be exposed to heavy metals in soil through:

1. Ingestion – swallowing contaminated soil (children at play, accidental consumption)
2. Inhalation – breathing in resuspended dust particles (dry season risk)
3. Dermal contact – skin exposure during agriculture or outdoor activities
4. Consumption of crops – long-term exposure via food chain (bioaccumulation)

Table 4.3 Toxicological Profiles of Key Metals

Element	Health Impact	Toxic Dose/Notes
Fe	Needed for hemoglobin, but excess may cause oxidative stress, especially in children	Toxic at >500 ppm (WHO limit: 425.5 ppm)
Cu	Excess leads to liver/kidney damage, vomiting, gastrointestinal distress	WHO limit: 36 ppm
Pb	Neurotoxic: affects IQ, memory, learning (especially in children); causes anemia, kidney damage	No safe exposure level; WHO limit: 85 ppm
Cr	Hexavalent Cr (Cr ⁶⁺) is carcinogenic , damages skin, liver, and respiratory system	WHO limit: 64 ppm
Mn	Overexposure linked to neurological issues (similar to Parkinson's disease)	WHO limit: 2000 ppm
Zn	Essential but high doses cause nausea, anemia, interference with Cu/Fe absorption	WHO limit: 300 ppm
Cd	Not detected; if present, causes kidney damage and bone demineralization	WHO limit: 3 ppm

Table 4.4 Risk Characterization: Kenana Study Area

Element	Average Study Area (ppm)	Risk Assessment
Cu	41.11 (dry), 38.56 (wet)	Above WHO limit — Chronic exposure through ingestion/inhalation may cause liver/kidney harm
Cr	82.57 (dry), 78.21 (wet)	Above WHO limit — Strong carcinogenic potential, especially for field workers and children
Pb	27.61 (dry), 25.79 (wet)	Below WHO limit, but still dangerous for children — neurotoxic at low doses
Fe	~22.5	Below threshold , no acute risk but should be monitored
Mn	~830	Well below limit , low risk but could contribute to cumulative neurotoxicity
Zn	~41	Safe , but elevated levels may affect crop safety long-term
Cd	ND	No direct risk detected , continue monitoring

5.4 Sensitive Populations at Risk

- Children (most vulnerable due to ingestion and developmental effects of Pb and Cr)
- Farmers and workers (dermal contact, inhalation)
- Pregnant women (risk to fetal development from Pb, Cr)
- Consumers of local crops (bioaccumulation of Cu, Cr)

5.5 Implications for Agriculture

- Soil-to-plant transfer of Cu and Cr may lead to contaminated food chains.
- Chronic exposure through vegetables and grains poses a long-term carcinogenic and toxicological risk.

Tabl 4.5 Summary of Human Health Risk by Metal

Metal	Acute Risk	Chronic Risk	Priority for Control
Cu	Moderate	High (long-term)	✓
Cr	High	Very High (carcinogenic)	✓✓✓
Pb	Low	High in children	✓✓
Fe	Low	Moderate (in accumulation)	✓
Mn	Low	Low–Moderate	△□
Zn	Low	Low	✗
Cd	None	None	✗

6. Conclusion

This study assessed heavy metal contamination in soils of the Kenana industrial area compared to a nearby control area, across both dry and wet seasons. Atomic Absorption Spectrophotometry (AAS) was used to measure the concentrations of Fe, Cu, Pb, Cr, Mn, and Zn. Results were evaluated against WHO permissible limits for soil quality.

Key conclusions include:

1. Copper (Cu) and Chromium (Cr) concentrations in the Kenana study area exceeded WHO safety limits in both seasons. This suggests ongoing industrial discharge or use of Cu/Cr-rich materials or chemicals.
2. Lead (Pb) levels, while below WHO limits, were significantly elevated in the study area (2× higher than control), posing particular risks to children and long-term health.
3. Iron (Fe), Manganese (Mn), and Zinc (Zn) were within WHO limits, but consistently higher in the study area, indicating pollution from industrial processes or agrochemical residues.
4. Cadmium (Cd) was not detected, which is a positive indicator for this region, reducing immediate concerns about this highly toxic metal.
5. Seasonal variation showed that dry seasons had generally higher concentrations of Cu, Cr, and Pb, likely due to reduced leaching and increased dust deposition. Wet seasons showed slight reductions, except for Fe and Mn, which rose due to increased mineral weathering.
6. Human health risk analysis showed high chronic risk from Cu and Cr, and moderate neurotoxic risk from Pb — particularly to children, farmers, and long-term residents.

This confirms that industrial activities in the Kenana area have significantly altered soil chemistry, creating localized but serious contamination patterns that may lead to health and ecological problems if left unmanaged.

Recommendations

Based on the findings, the following actions are recommended to reduce environmental and health risks:

7.1 Environmental Management

7.2 Health Protection

7.3 Policy and Research

7.4 Future Studies

Reference

- Navarro, M. C., Pérez-Sirvent, C., Martínez-Sánchez, M. J., Vidal, J., Tovar, P. J., & Bech, J. (2008). Abandoned mine sites as a source of contamination by heavy metals: A case study in a semi-arid zone. *Journal of Geochemical Exploration*, 96(2–3), 183–193.
2. Brumelis, G., Brown, D. H., Nikodemus, O., & Tjarve, D. (1999). The monitoring and risk assessment of Zn deposition around a metal smelter in Latvia. *Environmental Monitoring and Assessment*, 58(2), 201–212.
3. Vaalgamaa, S., & Conley, D. J. (2008). Detecting environmental change in estuaries: Nutrient and heavy metal distributions in sediment cores from the Gulf of Finland, Baltic Sea. *Estuarine, Coastal and Shelf Science*, 76(1), 45–56.
4. Cortés, O. E. J., Barbosa, L. A. D., & Kiperstok, A. (2003). Biological treatment of industrial liquid effluent in copper production industry. *Tecbahia Revista Baiana de Tecnologia*, 18(1), 89–99.
5. Devkota, B., & Schmidt, G. H. (2000). Accumulation of heavy metals in food plants and grasshoppers from the Taigetos Mountains, Greece. *Agriculture, Ecosystems & Environment*, 78(1), 85–91.
6. Baker, A. J. M. (1981). Accumulators and excluders: Strategies in response of plants to heavy metals. *Journal of Plant Nutrition*, 3, 643–654.
7. De Vries, W., Romkens, P. F., & Schütze, G. (2007). Critical soil concentrations of cadmium, lead, and mercury in view of health effects on humans and animals. *Reviews of Environmental Contamination and Toxicology*, 191, 91–130.
8. Galas-Gorchev, H. (1991). Dietary intake of pesticide residues: Cadmium, mercury, and lead. *Food Additives and Contaminants*, 8(6), 793–806.
9. IOCC, & CAOBISCO. (1996). *Heavy metals rapport*.
10. Wieting, J. (1988). Effects of air pollutants on groundwater quality in the Federal Republic of Germany. *Wasser Boden*, 40(4), 183–186.
11. Angelovičová, L., & Fazekašová, D. (2014). Contamination of the soil and water environment by heavy metals in the former mining area of Rudňany (Slovakia). *Soil & Water Research*, 9(1), 18–24.
12. Lee, C. S., Li, X., Shi, W., Cheung, S. C., & Thornton, I. (2006). A multivariate analysis of water quality in Lake Naivasha. *Science of the Total Environment*, 356(1), 45–54.
13. Skalski, T., Kędzior, R., Kolbe, D., & Knutelski, S. (2015). Ground beetles as indicators of heavy metal pollution in forests. *Sylwan*, 159(11), 905–911.
14. Alloway, B. J. (2012). *Heavy metals in soils: Trace metals and metalloids in soils and their bioavailability* (Vol. 22). Springer.

15. Castaldi, P., Melis, P., Silveti, M., et al. (2009). Influence of pea and wheat growth on Pb, Cd, and Zn mobility and soil biological status in a polluted amended soil. *Geoderma*, 151(3–4), 241–248.
16. Facchinelli, A., Sacchi, E., & Mallen, L. (2001). Multivariate statistical and GIS-based approach to identify heavy metal sources in soils. *Environmental Pollution*, 114(3), 313–324.
17. Fazekašová, D., Barančíková, G., Torma, S., Ivanová, M., & Manko, P. (2014). Chemical and environmental aspects of environmental and landscape components. In *5th Central European Conference in Regional Science (CERS)*.
18. Friedlová, M. (2010). The influence of heavy metals on soil biological and chemical properties. *Soil and Water Research*, 5(1), 21–27.
19. Huttmanová, E., Adamisin, P., & Hronec, O. (2015). Possibilities of soil revitalization in Slovakia towards sustainability. *European Journal of Sustainable Development*, 4(2), 121–128.
20. Kobza, J., & Gašová, K. (2014). Soil quality and heavy metal contamination. *Journal of Agricultural Science and Technology A*, 4(6), 504–513.
21. Senesil, G. S., Baldassarre, G., Senesi, N., & Radina, B. (1999). Trace element inputs into soils by anthropogenic activities and implications for human health. *Chemosphere*, 39(2), 343–377.
22. Takáč, P., Kozáková, L., Val'ková, M., & Zeleňák, F. (2008). Heavy metals in the middle Spiš soils. *Acta Montanistica Slovaca*, 13, 82–86.
23. Hronec, O., Tóth, J., & Tomáš, J. (2002). *Extraneous matter and their risks*. HarlequinQuality.
24. APHA. (2017). *Standard methods for the examination of water and wastewater* (23rd ed.). American Public Health Association.